

Abstract Submitted to the
International Conference on Strongly Correlated Electron Systems
University of Michigan, Ann Arbor
August 6-10, 2001

**High Resolution Bulk Sensitive Photoemission Study of the
Metal-Insulator Transition in $(V_{1-x}Cr_x)_2O_3$: Comparison to
LDA+DMFT Theory**

Hyeong-Do Kim¹, J. W. Allen¹, J.-H. Park², A. Sekiyama³, A. Yamasaki³, K. Kadono³, S. Suga³, Y. Saitoh⁴, T. Muro⁴, P. Metcalf⁵

¹ *University of Michigan, Ann Arbor, USA*

² *POSTECH, Pohang, Korea*

³ *Osaka University, Osaka, Japan*

⁴ *JAERI, Hyogo, Japan*

⁵ *Purdue University, West Lafayette, USA*

We present the first well resolved high photon energy bulk sensitive photoemission spectra for all phases of $(V_{1-x}Cr_x)_2O_3$ and compare them in detail to recent LDA+DMFT theory [1]. We find for the metal phase impressive general agreement that is sufficient to warrant quantitative optimization of the theory parameters, but for the paramagnetic insulating phase considerable differences which may not be fully explained by the high temperature of the calculations.

This work was supported by the U.S. NSF at U. Mich. (DMR-94-23741) and by the U.S. DoE at U. Mich. (DE-FG-02-90ER45416); and by Grant-in-Aid for COE Research (10CE2004) of MEXT, Japan and JASRI (No. 2000B0335-NS-np).

[1] K. Held *et al.*, Phys. Rev. Lett. **86**, 5345 (2001).